



WHO - GMP



AMOX LA AP

SUSPENSION FOR INJECTION, LONG ACTIVE

- **COMPOSITION:** Each 100 ml contains:
Amoxicillin (as trihydrate).....15 g
Excipient q.s.....100 ml
- **INDICATIONS:**
For the treatment of pneumonia, pasteurellosis, metritis, kidney - bladder inflammation, enteritis - diarrhea, edema disease caused by *E. coli*, *Salmonella*, wound infection, dermatitis in cattle, goats, sheep, pigs.
- **DOSAGE AND ADMINISTRATION:**
Intramuscular injection (I.M), single dose, repeat after 48 hours if necessary. Shake well before use:
- **Cattle, goats, sheep, pigs:** 1 ml/ 10 - 15 kg body weight.
- **Calves, kids, lambs:** 1 ml/ 10 kg body weight.
- **NOTE:**
- Shake well before use.
- Do not mix the medicine with any type on injection water.
- **CONTRAINDICATION:**
Do not use for animals in case of hypersensitivity to drug, or animals with impaired liver function.
- **WITHDRAWAL PERIOD:**
- **Meat:** 7 days.
- **Milk:** 4 days.
- **STORAGE RECOMMENDATION:**
Store in a cool and dry environment below 25°C.
Protect from light and extreme conditions.
- **MANUFACTURE:** SAI GON VET MANUFACTURING AND TRADING JOINT STOCK COMPANY



PACKING: 100 ml



SAI GON VET MANUFACTURING AND TRADING JOINT STOCK COMPANY

Anova Corporation Industrial Zone, Long Cang commune, Can Duoc district, Long An province, Vietnam
Email: info@saigonvet.vn - Tel: (+84 272) 372 62 62



WHO - GMP



CEPTIFUR AP

SUSPENSION FOR INJECTION
EFFECTIVE TREATMENT
FOR ACUTE PNEUMONIA, METRITIS

- **COMPOSITION:** Each 100 ml contains:
Ceptiofur (as HCl).....5 g
Excipient q.s.....100 ml
- **INDICATIONS:**
Ceptiofur is a broad-spectrum and new generation antibiotic. CEPTIFUR AP is used effectively for the treatment of pasteurellosis, acute pneumonia, salmonellosis, metritis and urinary tract infections in cattle, pigs, goats, sheep, dogs and cats.
- **DOSAGE AND ADMINISTRATION:**
Shake well before injection. For intramuscular or subcutaneous injection, once daily for 3 - 5 consecutive days:
- **Cattle:** 1 - 2 ml/ 50 kg body weight.
- **Pigs:** 1 - 3 ml/ 50 kg body weight.
- **Goats, sheep:** 1 ml/ 22 kg body weight.
- **Dogs, cats:** 1 ml/ 15 kg body weight.
- **CONTRAINDICATION:**
Do not use in case of hypersensitivity to antibiotics of cephalosporin and β - lactam group.
- **WITHDRAWAL PERIOD:**
- **Meat:** 5 days.
- **Milk:** 0 days.
- **STORAGE RECOMMENDATION:**
Store in a cool and dry environment below 25 °C.
Protect from light and extreme conditions.
- **MANUFACTURE:** SAI GON VET MANUFACTURING AND TRADING JOINT STOCK COMPANY



PACKING: 100 ml



SAI GON VET MANUFACTURING AND TRADING JOINT STOCK COMPANY

Anova Corporation Industrial Zone, Long Cang commune,
Can Duoc district, Long An province, Vietnam
Email: info@saigonvet.vn - Tel: (+84 272) 372 62 62